

Differential expression of miR200a-3p and miR21 in grade II–III and grade IV gliomas

Evidence that miR200a-3p is regulated by O⁶-methylguanine methyltransferase and promotes temozolomide responsiveness

Yolande Berthois^{1,*}, Christine Delfino¹, Philippe Metellus^{1,2}, Frederic Fina³, Isabelle Nanni-Metellus³, Hayat Al Aswy¹, Victor Pirisi¹, L'Houcine Ouafik^{1,3}, and Françoise Boudouresque¹

¹Aix Marseille Université; Inserm; CRO2 UMR_S 911; Marseille, France; ²Département de Neuropathologie; APHM; Hopital Timone; Marseille, France;

³Service de Transfert d'Oncologie Biologique; APHM; Hopital Nord; Marseille, France

Keywords: glioblastoma, microRNA, miR-200a-3p, miR21, temozolomide, O⁶-methylguanine methyltransferase, chemoresistance

Abbreviations: EMT, epithelial–mesenchymal transition; FBS, bovine serum; GBM, glioblastoma; GFAP, glial fibrillary acidic protein; LG, lower grade; MGMT, O⁶-methylguanine methyltransferase; miRNA, microRNA; MMLV, moloney murine leukemia virus reverse transcriptase; MTT, thiazolyl blue tetrazolium blue; O⁶-BG, O⁶-benzylguanine; PI3K, phosphoinositide-3-kinase; PTEN, phosphatase and tensin homolog; TMZ, temozolomide; RT-PCR, reverse transcription polymerase chain reaction; STUPP, radiotherapy plus concomitant and adjuvant temozolomide; WHO, World Health Organization

Glioblastoma multiforme (GBM) is the most common primary brain tumor and is among the deadliest of human cancers. Dysregulation of microRNAs (miRNAs) expression is an important step in tumor progression as miRNAs can act as tumor suppressors or oncogenes and may affect cell sensitivity to chemotherapy. Whereas the oncogenic miR21 has been shown to be overexpressed in gliomas, the expression and function of the tumor-suppressor miR200a in GBMs remains unknown. In this study, we show that miR21 is upregulated in grade IV (GBMs) vs. grade II–III (LGs) gliomas, confirming that miR21 expression level is correlated with tumor grade, and that it may be considered as a marker of tumor progression. Conversely, miR200a is demonstrated for the first time to be downregulated in GBMs compared with LGs, and overexpression of miR200a in GBM cells is shown to promote TMZ-sensitivity. Interestingly, miR200a but not miR21 expression level is significantly higher in TMZ-responsive vs. -unresponsive tumoral glial cells in primary culture. Furthermore, miR200a appears negatively correlated with the expression of the DNA repair enzyme O⁶-methylguanine methyltransferase (MGMT), and the inhibition of MGMT activity results in an increase of miR200a expression in GBM cells. Taken together, these data strongly suggest that miR200a is likely to act as a crucial antitumoral factor regarding glioma progression. Interplay between miR200a and MGMT should be considered as potential mechanism involved in therapeutic response.

Introduction

Glioblastoma multiforme (GBM) is the most common primary brain tumor and among the deadliest of human cancers. Despite advances in diagnosis and treatment, mean survival time does not exceed 2 y. These tumors typically display a very high proliferative rate with widespread microvascular development and areas of focal necrosis.¹ The current standard therapy for GBM includes surgery with maximum feasible resection, radiotherapy, and treatment with the monofunctional alkylating agent temozolomide (TMZ).^{2,3} Although TMZ has improved overall and progression-free survival, a majority of patients still

experience disease progression within 1 y. Indeed, GBMs can present innate resistance or develop acquired resistance during TMZ treatment.^{4–6}

Resistance of GBMs to TMZ is reported to be mainly, but not exclusively, dependent on high levels of the DNA repair enzyme, O⁶-methylguanine methyltransferase (MGMT), which can reverse the methylation damage induced by alkylating agents.^{7–9} Although a number of studies have shown that a deficiency of MGMT can augment the sensitivity of GBMs to alkylating agents such as TMZ, a panel of tumors with low levels of MGMT are nevertheless chemoresistant.^{10,11} This suggests that additional mechanisms are involved in tumor resistance to

*Correspondence to: Yolande Berthois; Email: yolande.berthois@univ-amu.fr
Submitted: 01/21/2014; Revised: 04/16/2014; Accepted: 04/16/2014
<http://dx.doi.org/10.4161/cbt.28920>

chemotherapy.¹² Thus, resistance of GBM cells to TMZ includes various events, such as loss of the phosphatase and tensin homolog (PTEN), leading to the activation of the phosphoinositide-3-kinase (PI3K)/Akt pathway, deficiency in the DNA mismatch repair system, mutations in the pro-apoptotic p53 protein, over-expression of the anti-apoptotic protein Bcl-2, or the selection of less-differentiated pre-existing resistant cells in the parental tumor.¹³⁻¹⁹

Dysregulation of the expression of microRNAs (miRNAs) is emerging as an important step in early processes during tumorigenesis as well as disease progression/metastasis in various malignancies including GBMs.²⁰⁻²² The miRNAs function mainly through their ability to bind to the 3' untranslated regions of mRNAs, leading to their degradation or suppression of their translation.²³ MiRNAs can act as tumor suppressors or oncogenes depending upon the nature of the targeted genes.²⁴ MiRNA profiles have been used successfully to classify tumors and predict response to therapies.^{25,26} These analyses have identified a number of miRNAs consistently altered in gliomas, and one predominant miRNA to arise from these studies is miR21.²⁷ Thus, the miRNA expression studies demonstrated that miR21 is overexpressed in gliomas and glioma cell lines relative to normal tissue.²⁸ However, the association of miR21 with tumor grade is less consistent. While some studies indicate that miR21 is upregulated in both low- and high-grade gliomas, other suggest that it is associated with progression and upregulated more frequently in high-grade tumors.^{28,29} Whereas the correlation of miR21 with clinical outcome in gliomas is little documented, miR21 expression level has been correlated with overall and disease-free survival and suggested to be a biomarker for chemoresistance in other types of cancer including leukemia, pancreatic and lung cancers.³⁰⁻³³

Besides oncogenic miRNAs such as miR21, a number of miRNAs have been described to elicit tumor suppressive role and to display downregulation during tumor development.²⁴ This is the case for the miR200 family, composed of five members: miR-200a, -200b, -200c, -141, and -429. Members of the miR200 family have been shown to be major regulators of the EMT (epithelial-mesenchymal transition) process, primarily by targeting EMT-driving transcription factors ZEB1 and ZEB2 (SIP1), which in turn transcriptionally repress miR200 family members in a double negative feedback loop.^{34,35} A number of studies indicate that members of the miR200 family are down-regulated in a variety of human cancers such as breast, ovarian and lung cancers.³⁶⁻³⁸ Moreover, a role for miR200 family in drug resistance has been reported in several cancers. Thus, gemcitabine-resistant pancreatic cancer is associated with decreased miR200 expression.³⁹ Similarly, expression level of miR200a is correlated with resistance of ovarian and bladder cancers to paclitaxel and anti-EGFR therapy, respectively.^{36,40} However, contradictory data regarding these microRNAs have been reported, since other studies describe overexpression of miR200 family members in cancers such as melanoma, ovarian and colorectal cancers.⁴¹⁻⁴⁴ Contrary to miR21, the role of miR200 in brain tumors is little documented. Whereas Saydam et al.⁴⁵ have shown that down-regulation of miR200a in meningiomas increased tumor growth through the activation of the wnt/ β -catenin signaling pathway,




Figure 1. Effect of temozolomide (TMZ) on the proliferation of tumoral glial cell lines (A) and glioblastoma-derived cell primary cultures (B). Cells were treated for 3 d in the presence of the indicated concentrations of TMZ. Cell proliferation was then assessed by MTT metabolism. Values are expressed as % of untreated control cells. Data are the mean \pm SD of 3 separate experiments.

the expression and function of miR200 in glioblastomas have not been investigated and remain unknown.

In order to clarify the functional role of miR21 and more specifically miR200a in glioblastomas, the expression level of both miRNAs was first analyzed and compared in grade IV (GBMs) and grade II–III (LGs) glioma samples. We show that miR21 is upregulated in GBMs when compared with LGs, confirming that miR21 expression level is correlated with tumor grade, and that it may be considered as a marker of tumor progression. Conversely, miR200a is demonstrated for the first time to be down-regulated in GBMs compared with lower grade gliomas. Interestingly, downregulation of miR200a in GBMs is correlated with MGMT expression level and TMZ-resistance, strongly suggesting that this miRNA is likely a crucial factor when considering glioma progression and therapeutic response.

Results

Effect of TMZ on glioma cell proliferation

To evaluate the antitumor effect of TMZ in malignant glioma cells, several glial tumoral cell lines were treated with increased doses of TMZ for 72 h. Cell viability was then assessed by MTT metabolism. As shown in Figure 1A, variable cell growth inhibitory effects of TMZ were observed, depending on the tumor cell lines examined. Whereas U138, T98G, and LN18 cells were

Table 1. Characterization of the tumoral glial cell lines used in this study

Cell lines	MGMT/18s	MGMT protein	MGMT methylation	miR200a/U6	TMZ IC ₅₀ μ M
U87	1.0 10 ⁻⁶ \pm 2.010 ⁻⁶ [1]*	nd	M	2.0 10 ⁻³ \pm 1.1 10 ⁻³	50
U373	5.1 10 ⁻⁶ \pm 2.8 10 ⁻⁶ [5.1]	nd	M	1.1 10 ⁻³ \pm 2.2 10 ⁻³	110
T98G	1.1 10 ⁻³ \pm 0.9 10 ⁻³ [1.1 10 ³]	+	M	1.0 10 ⁻⁴ \pm 0.07 10 ⁻⁴	>300
U118	2.0 10 ⁻³ \pm 1.1 10 ⁻³ [2 10 ³]	+	UM	1.0 10 ⁻³ \pm 0.6 10 ⁻⁴	220
U138	2.4 10 ⁻² \pm 0.6 10 ⁻² [2.4 10 ⁴]	++	M+UM	1.2 10 ⁻⁴ \pm 1.1 10 ⁻⁴	>300
LN18	3.1 10 ⁻² \pm 0.3 10 ⁻² [3.1 10 ⁴]	++	UM	3.1 10 ⁻⁴ \pm 1.9 10 ⁻⁴	>300

*in brackets, relative expression level of MGMT, taking MGMT level in U87 cells as 1; nd, no detectable; M, methylated MGMT; UM, unmethylated MGMT.

Table 2. Characterization of the glioblastoma-derived cell primary cultures used in this study

Primary cultures	MGMT/18s	MGMT protein	MGMT methylation	miR200a/U6	TMZ IC ₅₀ μ M
GBM.4	0.21 \pm 0.08 [2 10 ⁵] *	+	M	1.1 10 ⁻² \pm 1.6 10 ⁻²	>300
GBM.5	7.0 10 ⁻³ \pm 0.8 10 ⁻³ [5 10 ³]	+	M	0.11 \pm 0.51	>300
GBM.7	2.5 10 ⁻³ \pm 0.7 10 ⁻³ [2.5 10 ³]	+	UM	0.53	>300
GBM.8	5.0 10 ⁻³ \pm 1.1 10 ⁻³ [5 10 ³]	+/-	M		>300
GBM.9	2.1 10 ⁻⁴ \pm 1.0 10 ⁻⁴ [2 10 ²]	nd	M	1.12 \pm 0.22	250
GBM.10	0.9 10 ⁻⁴ \pm 0.3 10 ⁻⁴ [0.9 10 ²]	nd	M		250
GBM.11	2.3 10 ⁻⁴ \pm 0.9 10 ⁻⁴ [2.3 10 ²]	nd	M	0.12	220
GBM.12	0.83 \pm 0.03 [8 10 ⁵]	++	UM	3.0 10 ⁻³ \pm 0.6 10 ⁻³	>300
GBM.14	7.0 10 ⁻³ \pm 2.3 10 ⁻² [7 10 ³]		UM	0.4 \pm 2.1 10 ⁻²	220
GBM.15	0.80 \pm 0.05 [8 10 ⁵]	++	UM	1.33 10 ⁻³ \pm 0.11 10 ⁻³	>300
GBM.16	5.0 10 ⁻² \pm 0.11 [5 10 ⁴]	++	UM	1.50 10 ⁻² \pm 0.07 10 ⁻²	220
GBM.19	2.3 10 ⁻³ \pm 0.6 10 ⁻³ [2.3 10 ³]		UM	1.9 10 ⁻² \pm 0.7 10 ⁻³	>300
GBM.21	3.0 10 ⁻³ \pm 0.9 10 ⁻⁴ [3 10 ³]	++	UM	6.8 10 ⁻² \pm 0.8 10 ⁻²	>300
GBM.22	1.8 10 ⁻² \pm 1.0 10 ⁻⁴ [1.8 10 ⁴]	++	UM	7.3 10 ⁻³ \pm 1.6 10 ⁻³	>300
GBM.23	0.12 \pm 0.05 [1.2 10 ⁵]	++	UM		>300
GBM.24	1.9 10 ⁻³ \pm 1.3 10 ⁻³ [1.9 10 ³]		M	1.5 10 ⁻² \pm 2.1 10 ⁻²	>300
GBM.25	2.0 10 ⁻⁶ \pm 0.3 10 ⁻⁶ [2]	nd	M	1.64 \pm 0.11	180

*in brackets, relative expression level of MGMT, taking MGMT level in U87 cells as 1 (cf **Table 1**); nd, no detectable; M, methylated MGMT; UM, unmethylated MGMT.

found to be little affected by TMZ (IC₅₀ > 300 μ M), a dose-dependent inhibition of the proliferation of U87, U373, and U118 cells was observed, with IC₅₀ of 50 μ M, 110 μ M, and 220 μ M respectively. The effect of TMZ on cell proliferation was also examined in a panel of human tumoral glial cells established in primary culture from glioblastomas. Data reported in **Figure 1B** indicate that most of the primary cultures were unaffected by the presence of TMZ (150–300 μ M). However, a dose-dependent inhibition of cell proliferation was observed in two primary cultures treated with TMZ (GBM.14 and GBM.16), with IC₅₀ of 220 μ M. The IC₅₀ values of TMZ for each cell line and glioblastoma primary culture are listed in **Tables 1 and 2**, respectively.

In order to verify the mechanism through which TMZ affects cell glioma proliferation, its effect on cell cycle was further analyzed. In agreement with a number of reports, our data confirm that the growth-inhibitory effect of TMZ is associated with accumulation of cells into G₂/M cell cycle phases and concomitant

decrease of cell number into G₀/G₁ (**Figs. S1 and S2**).⁴⁶⁻⁴⁸ These results confirm that TMZ is able to affect glioma cell growth by blocking the transit beyond cell cycle phases.

Expression level of molecular markers in GBM and lower grade gliomas tissue samples

A number of genes are associated with tumor grade, progression and patient survival with GBM.^{46,47,49,50} The expression of several genes of interest was measured in a panel of high (GBM) and lower grade gliomas. In parallel, several tumoral cell lines issued from GBM were included in the study. Data reported in **Figure 2** indicate that the expression of PDGFR- α , VEGF α , BCAN, MGMT, and Bcl₂:Bax ratio were not significantly different in GBMs, tumoral cell lines, and lower grade gliomas. On the contrary, TIMP1, YKL-40 (chitinase), and vimentin, described to be implicated in the malignancy of gliomas, appeared over-expressed in GBMs and tumoral cell lines when compared with lower grade gliomas.




Figure 2. Relative expression of biomarkers in grade IV (GBMs) and grade II–III gliomas (low grade) and in tumoral glial cell lines. Expression levels were measured by quantitative RT-PCR and values were normalized with 18s housekeeping gene. For each data set, median value and 5% and 95% confidence intervals are reported. ** $P < 0.01$; *** $P < 0.001$




Figure 3. Relative expression of miR-21 (**A and C**) and miR200a (**B and D**) in grade IV (GBM) and grade II–III gliomas (low grade), and in tumoral glial cell lines. Expression levels were measured by quantitative RT-PCR and values were normalized with snU6 housekeeping gene. For each data set, median value and 5% and 95% confidence intervals are reported (**C and D**). ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$.

MiR200a and miR21 expression levels in GBM tissue and lower grade gliomas tissue samples

In order to determine whether miR21 and miR200a may be associated to brain tumor progression, the expression of both miRNAs was measured in lower grade (LG) and GBM samples, as well as in tumoral glial cell lines. The relative expression levels reported in **Figure 3** demonstrate that miR21 was significantly overexpressed in GBM compared with LG samples ($P < 0.001$). Similar results were observed when comparing miR21 expression in GBM-derived cell lines vs. LG samples ($P < 0.01$). Conversely, miR200a expression level was significantly lower in both GBM samples and glioma cell lines, compared with LG ($P < 0.0001$ and $P < 0.001$, respectively).

miR200a and miR21 affect TMZ-tumoral glial cell responsiveness

An increasing number of microRNAs are described to be involved in tumoral cell response to chemotherapy. To evaluate the potential role of miR200a and miR21 in tumoral glial cell responsiveness to TMZ, the expression of both miRNAs was first determined in a panel of glioblastoma cell lines (U87, U373, T98, and LN18) treated in the absence or in the presence of TMZ. Our data indicate that under our experimental conditions, TMZ was unable to affect the expression of miR200a and miR21 in all the cell lines as well as in several GBM primary cultures (GBM.10, GBM.11, GBM.14, and GBM.16) examined (not shown).

In a second time, miRNA transfection studies were performed to assess the potential involvement of miRNA, in particular miR200a, in TMZ-responsiveness. For this purpose, U87 and U373 cell lines were transfected with either the antagomir of miR21 (as-miR21) or the mimic pre-miR200a. Control cells were transfected with the respective control oligonucleotides. RT-PCR measurements performed 24 h and 48 h post-transfection demonstrated decreased expression of miR21 and overexpression of miR200a, respectively (Figs. S3 and S4). Whereas TMZ at the indicated concentrations had no significant effect on the proliferation of cells transfected with pre-miRNA control, the addition of TMZ in U87 and U373 cells overexpressing miR200a decreased cell growth to 60% of control (Fig. 4). As expected, similar results were obtained after cell transfection with the miR21 inhibitor. Furthermore, the effect of miRNA transfection on TMZ-responsiveness was verified in two GBM primary cultures, previously shown to display different growth response to TMZ. Thus, whereas the proliferation of GBM.16 primary culture was reduced in the presence of TMZ ($IC_{50} = 220 \mu M$), GBM.22 appeared to be TMZ-unresponsive (cf. Fig. 1). Data reported in Figure 4 show that TMZ decreased the proliferation of GBM.16 overexpressing miR200a to 50% of control, *vs* 82% in cells transfected with pre-miRNA control. Likely, the reduction of miR21 expression in the same cells resulted in an increased TMZ growth inhibition. Conversely, neither miR200a overexpression nor miR21 inhibition could modify the response of GBM.22 primary culture to TMZ.

miR200a expression is correlated with MGMT expression and TMZ glial cell responsiveness

Since miR200a and miR21 were shown to be involved in TMZ-responsiveness of a number of tumoral glial cells, data analysis was conducted to verify whether both miRNAs could be used to discriminate TMZ-responsive and -unresponsive GBMs. GBM primary cultures were then divided into two subgroups. The first one contained primary cultures that displayed growth inhibition in the presence of TMZ ($IC_{50} = 200-250 \mu M$). The second group included primary cultures that were unaffected by the presence of TMZ in our experimental conditions ($IC_{50} > 300 \mu M$). Data relative to the characterization of each primary culture used in this analysis are reported in Table 2.

As MGMT is largely admitted to play a crucial role in mediating the resistance to alkylating agents such as TMZ, the correlation between MGMT expression and TMZ responsiveness was examined. Immunoblot analysis of MGMT in a panel of GBM primary cultures verified that MGMT expression was associated, in most of them, with the presence of MGMT protein (Fig. 5; Table 2). Data reported in Figure 6A confirm that MGMT expression levels were significantly higher in GBM primary




Figure 4. Effect of miR21 inhibition and miR200a overexpression on TMZ-responsiveness. Tumoral glial cell lines (U87, U373) and glioblastoma-derived primary cultures (GBM.16, GBM.22) were transfected with anti-sens miR21 (as-miR21) or mimic premature miR200a (pre-miR200a). Control cells were transfected with anti-sens miRNA control (as-miR control) or premature miRNA control (pre-miR control). Three days post-transfection, cells were treated for 3 d with the indicated concentrations of TMZ. Cell proliferation was then assessed by MTT metabolism. Values are expressed as % of the respective untreated control cells. Data are the mean \pm SD of triplicate and are representative of 2 separate experiments.




Figure 5. Expression of MGMT protein in GBM-derived primary cultures and tumoral glial cell lines. Cell lysates were prepared as described in Materials and Methods. After separation by electrophoresis and transfer on nitrocellulose membrane, MGMT protein was analyzed by immunoblotting. In parallel, β-actin protein level in each sample was verified. Data are representative of two separate experiments.




Figure 6. Correlation analysis between miR200a and MGMT expression levels and TMZ-responsiveness in GBM-derived cell primary cultures (A–C) and tumoral glial cell lines (D–F). Primary cultures and tumoral cell lines were subdivided into TMZ-responsive ($IC_{50} = 200$ – 250 μ M and 50 – 220 μ M, respectively) and -unresponsive ($IC_{50} > 300$ μ M) groups. For each groups, median values and 5% and 95% confidence intervals for MGMT (A and D) and miR200a (B and E) are reported. Correlation analysis between MGMT and miR200a expression levels are presented (C and F).

cultures resistant to TMZ, compared with TMZ-responsive group ($P = 0.042$). Whereas both subgroups were shown to express similar levels of miR21 (not shown), miR200a expression was significantly increased in TMZ-responsive GBM cultures ($P = 0.030$) (Fig. 6B). Given these data, a potential correlation between miR200a and MGMT expressions was then examined. As expected, a negative correlation was observable between miR200a and MGMT ($r^2 = -0.65$, $P = 0.019$) (Fig. 6C).

The same analysis was conducted on tumoral cell lines. The TMZ-responsive ($IC_{50} = 50$ – 220 μ M) and -unresponsive ($IC_{50} > 300$ μ M) groups included U87, U373, U118, and LN18, T98G, U138 cell lines, respectively. Whereas the expression of MGMT tended to be higher in TMZ-resistant cells (Fig. 6D), miR200a expression level was augmented in TMZ-sensitive cell lines (Fig. 6E). However, these differences in expression levels were not significant, probably due to the small number of cell lines examined. Nevertheless, tendency to a negative correlation between miR200a and MGMT expressions was observable ($r^2 = -0.73$, $P = 0.08$) (Fig. 6F).

miRNA200a expression is regulated by MGMT activity

The correlation observed between MGMT and miR200a expression levels in tumoral glial cells suggests that molecular interactions between MGMT and miR200a might occur. In a first time, experiment was conducted to verify that the inhibition of MGMT could promote the growth inhibitory effect of TMZ in tumoral glial cells. The TMZ-unresponsive LN18 and T98G cell lines that were shown to express significant levels of MGMT

(Fig. 5; Table 1) were treated with the MGMT inhibitor O⁶-BG (50 μ M) in the absence or in the presence of different concentrations of TMZ (100–300 μ M). Our results demonstrate that TMZ at 300 μ M was able to reduce T98G and LN18 cell proliferation to 66% and 60% of control respectively, when added in the presence of O⁶-BG (Fig. 7A; Fig. S5). In parallel, the disappearance of detectable MGMT protein level in both cell lines under O⁶-BG treatment was verified by western immunoblotting (Fig. 7B).

In order to determine whether the expression of miR200a and miR21 was dependent on MGMT activity, miRNA expression levels were then measured in cells treated with O⁶-BG. The inhibition of MGMT activity allowed to an important increase of miR200a expression in LN18 cells, 24 h after the addition of 20–50 μ M O⁶-BG (Fig. 8). A moderate effect persisted 72 h after treatment with 40–50 μ M O⁶-BG (not shown). Similarly, miR200a expression level appeared significantly augmented 24 h after the addition of 20–50 μ M O⁶-BG in T98G cells. Conversely, O⁶-BG was unable to significantly modify the expression of miR21 in both LN18 and T98G cell lines.

In a second step, potential regulation of MGMT expression by miRNA was examined. LN18 and T98G cells were then transfected with miR21 antagomir or pre-miR200a mimic, then MGMT expression was determined 48 h post-transfection. Results obtained demonstrate that neither reduction of miR21 nor overexpression of miR200a could modify the expression level of MGMT (not shown).

Discussion

It is now well established that particular miRNAs play a crucial role in gene regulation in health and disease. Notably, deregulation of a number of miRNAs has been demonstrated to be associated with cancer pathogenesis.²⁴ Furthermore, accumulating evidence has revealed that miRNAs may play a role in promoting tumoral cell response to chemotherapeutic agents, and a number of reports have proposed the use of miRNAs as potential predictive and prognostic factors in a variety of cancers including GBMs.^{29,51-53} In this study, we demonstrate differential expression of miR21 and miR200a in grade IV (GBM) vs. lower grade (II–III) (LG) human brain tumors. Moreover, our data reveal for the first time a specific function for miR200a in promoting the response of tumoral glial cells to the alkylating agent TMZ.

MiR21 has been described to be significantly augmented in GBMs and to function as an oncogene that targets a variety of tumor suppressor genes.^{27,30} However, the association of miR21 with grade in gliomas is less consistent. In agreement with the report by Hermansen et al.²⁹ we demonstrate herein that miR21 is overexpressed in GBM samples when compared with LGs. Likewise, cell lines derived from glioblastomas are shown to express higher miR21 levels than LGs, confirming that miR21 expression level is correlated with disease malignity, and indicating that it may discriminate grade IV from grade II–III gliomas. Contrary to miR21, the expression and function of miR200 family members in brain tumors are little documented. Saydam et al.⁴⁵ have shown that downregulation of miR200a results in the activation of the wnt/ β -catenin signaling pathway and increases tumor growth in meningiomas. Also, miR200 family has been demonstrated to be involved in the suppression of EMT and self-renewal of GBM initiating cells by Erismodegib.⁵⁴ In agreement with these data, we were able to demonstrate for the first time that miR200a is downregulated in both GBMs and tumoral glial cell lines when compared with LGs, strongly suggesting that loss or reduction of miR200a expression occurs during GBM development and tumor progression.

Innate or acquired resistance to chemotherapeutic agents such as TMZ is a common fate in GBMs.⁴⁻⁶ Consistent with this observation, we show that a few number of GBM primary cultures display growth inhibition in response to TMZ. Likewise, among 6 GBM-derived cell lines examined, 3 are shown to be TMZ-responsive. The presence of MGMT in tumoral glial cells has been shown to be responsible, at least in part, of TMZ resistance.^{7,8,55} This is comforted by the association that we observe, between MGMT expression level and TMZ-growth inhibitory activity in both tumoral cell lines and GBM primary cultures. However, published data indicate that this correlation is far from being total, arguing for the involvement of




Figure 7. Effect of the MGMT inhibitor O⁶-BG on TMZ-responsiveness of tumoral glial cells. **(A)** T98G and LN18 cell lines were pre-treated with 50 μM O⁶-BG for 4 h. Cells were then treated in the absence or in the presence of TMZ at the indicated concentrations. Three days later, cell proliferation was assessed by MTT metabolism. Values are expressed as % of the respective untreated control cells. Data are the mean ± SD of triplicate and are representative of 3 separate experiments. ***P* < 0.001. **(B)** the effect of different concentrations (10–50 μM) O⁶-BG on MGMT protein level was examined by immunoblotting. In parallel, β-actin expression was verified for each sample. Data are representative of two separate experiments.




Figure 8. Effect of the MGMT inhibitor O⁶-BG on miR200a expression level in tumoral glial cells. T98G and LN18 cell lines were treated for 24 h with different concentrations (10–50 μM) of O⁶-BG. Expression levels of miR200a were measured by quantitative RT-PCR and values were normalized with snU6 housekeeping gene. Data are expressed as % of untreated control cells. Values are the mean ± SD of triplicate and are representative of 3 separate experiments.

MGMT-independent mechanisms in cell resistance to TMZ.¹² Thus, besides their involvement in tumor development and progression, a number of miRNAs have been shown to influence cell responsiveness to chemotherapies.⁵⁶ This is true for miR21, as increasing number of studies demonstrate that knockdown of miR21 is able to promote tumor growth inhibition triggered by chemotherapeutic agents.^{57,58} Thus, miR21 has been described to mediate resistance to chemotherapy and ionizing radiation in GBMs through the inhibition of apoptotic signaling pathway.^{59,60} In agreement with these data we confirm that depletion of miR21 increases TMZ-responsiveness of tumoral glial U87 and U373 cell lines. On contrary, TMZ is unable to modify the expression levels of miR21 in several tumoral glial cell lines and GBM primary cultures. These late data are in apparent contradiction with the report by Wong et al.⁶¹ demonstrating that exposure of glioblastoma cells to TMZ elevated miR21 expression. However, miR21 augmentation was observed in glioblastoma cells that developed TMZ resistance after long-term, chronic administration of TMZ. In our study TMZ treatments did not exceed 72 h, suggesting that regulation of miR21 by TMZ occurs via indirect mechanisms and requires profound phenotypic and/or genotypic changes.

Although the role of miR200a in tumoral cell responsiveness to therapy is less documented, it is obvious that the expression of miR200a in a variety of tumoral cells promotes cytotoxic activity of antitumoral drugs.^{36,39,62} In agreement with these data, we show that the overexpression of miR200a in U87 and U373 cells increases the growth inhibitory activity of TMZ. Likely, increasing expression levels of miR200a promotes the response of GBM.16 primary culture to TMZ. Moreover, our results bring in evidence a correlation between miR200a expression level and in vitro growth response of GBM primary cultures to TMZ. Thus, primary cultures that display growth resistance to 300 μ M TMZ are shown to express lower miR200a levels than TMZ-responsive GBMs. Although TMZ-responsive cell lines tended to express higher miR200a levels when compared with unresponsive cells, significant correlation cannot be evidenced, possibly due to the small number of cell lines examined. Interestingly, our data further indicate a negative correlation between miR200a and MGMT expression levels in GBM primary cultures. Similarly, miR200a tended to be negatively correlated with MGMT in tumoral cell lines, suggesting that reciprocal regulation between miR200a and MGMT occurs in glioblastoma cells. In agreement with these data, the inhibition of MGMT activity by O⁶-BG in two different cell lines (LN18 and T98G), increases the expression of miR200a and partially reverses cell resistance to TMZ. However, modifying miR200a expression in LN18 and T98 cells did not affect MGMT level, suggesting unilateral regulation of miR200a by MGMT. The mechanism through which MGMT affects miR200a expression is still unknown. However, in deficient MGMT cells, O⁶-MeG adducts produced under the action of a variety of mutagens may activate apoptotic pathway in a p53-dependent manner.^{63,64} Furthermore, p53 has been shown to upregulate miR200 family members that in return suppress EMT by repressing the expression of ZEB1 and ZEB2 (SIP1).⁶⁵ Thus, we suggest that accumulation of O⁶-MeG adducts in cells

treated with O⁶-BG might induce miR200a expression through a p53-dependent mechanism. Certainly, future investigations are needed to define the functional relationship between decrease of miR200a expression, MGMT activity and oncogenic features.

In conclusion, our present data confirm overexpression of miR21 and demonstrate downregulation of miR200a in high grade (IV) gliomas. Importantly, miR200a is shown to be able to promote chemotherapy responsiveness when overexpressed in glioblastoma cells. Further, the demonstration of the upregulation of miR200a by inhibition of MGMT activity provides new insight into the mechanisms responsible of glioma cell progression and chemo-resistance.

Materials and Methods

Materials

Cell culture medium and reagents, oligofectamine, random hexamers, MMLV, mirVanaTM kit, turbo DNA-free kit, miRNA assay kit, miRNA reverse transcription kit, and PCR TaqMan probe assays were provided by Invitrogen Life Technologies Inc. Temozolomide, O⁶-benzylguanine (O⁶-BG), thiazolyl blue tetrazolium blue (MTT), and collagenase were from Sigma-Aldrich. Enhanced chemiluminescent substrate was from GE Healthcare Life Sciences. Protease inhibitor cocktail and standard genomic DNA was furnished by Roche Applied Science. Tri-Reagent and DNA purification kit were from Euromedex and Promega, respectively. Anti-MGMT monoclonal antibody (clone MT3.1) and horseradish peroxidase-conjugated secondary antibody were from Epitomics and Santa Cruz Biotechnology (Tebu) respectively. Epitect bisulfite kit was provided by Qiagen.

Cell line culture

Human glioblastoma cell lines U87, U373, T98G, LN18, and U138 were obtained from the American Type Culture Collection. They were maintained in DMEM supplemented with 2 mM glutamine and 10% fetal bovine serum (FBS), except LN18 cells that were cultured in the presence of 5% FBS. Cells were cultured at 37 °C, under a humid atmosphere of 5% CO₂/95% air.

Primary cultures of tumor glial cells

Tumor samples classified as glioblastoma based on World Health Organization (WHO) criteria were obtained from patients undergoing surgical treatment at the Department of Neurosurgery, CHU Timone. Within 1–4 h after surgical removal, tumors were washed and dissected into small pieces (<1 mm³) in RPMI-1640 medium supplemented with 2% FBS and 50 μ g/mL penicillin/streptomycin, and disrupted mechanically. Tumoral glial cells were separated on a 30% continuous dextran gradient prepared in RPMI, washed in RPMI and subsequently dissociated in 10 μ g/mL collagenase in RPMI. After 1 h at 37 °C, cells were washed before plating in T-75 cm² flasks. GBM primary cultures were maintained in DMEM/F12 supplemented with 10% FBS and 50 μ g/mL penicillin/streptomycin, at 37 °C in a humidified atmosphere of 5% CO₂/95% air. Cells were grown to confluence with medium change twice a week and used until passage 5. Glial cell cultures were characterized by the presence of GFAP and the absence of oligodendrocyte markers.

Patients

Molecular analysis was performed on 24 primary glioblastomas (GBMs) and 10 low grade brain tumors (LGs) resected at the Department of Neurosurgery, CHU Timone. All the tissue procurement protocols were approved by the relevant institutional committees (Aix-Marseille University) and were undertaken under informed consent of each patient or relatives. Diagnosis and grading of all tumors were certified by independent neuropathological examination. The histological diagnosis of GBM and LG was performed according to WHO guidelines. The mean age of the patients was 58.4 y (range = 27–81 y). Twenty-seven (79.4%) patients were males, and seven (20.6%) were females (Table 3). One viable sample from the tumors was stored at -80°C for molecular analysis and one was fixed in buffered-formalin and embedded in paraffin. For each sample, at least 1 slice was stained with hematoxylin and eosin to control the percentage of tumor cells. Only samples containing at least 60% of tumor cells were retained for subsequent analysis.

Preparation of cell extracts and western immunoblotting

Cells were seeded at 15000 cells/cm² in their regular culture medium. At pre-confluence, cells were rinsed with phosphate-buffered saline (PBS) and treated in fresh culture medium as mentioned. At the end of the treatment, cell monolayers were washed with ice-cold PBS and incubated in lysis buffer (25 mM TRIS-HCl pH 7.4, 150 mM NaCl, 2.5 mM EDTA, 50 mM sodium fluoride, 2 mM sodium orthovanadate, 1% NP40, 0.1% SDS, 0.5% sodium deoxycholate, 1 mM phenylmethylsulfonyl-fluoride) containing protease inhibitor cocktail as indicated by the manufacturer. Insoluble material was removed by centrifugation at 4°C for 15 min at $20000 \times g$. Cell extracts were resolved by SDS-PAGE and separated proteins were then electroblotted on a nitrocellulose membrane. Western immunoblotting was performed as previously described.⁶⁶

Quantitative real-time polymerase chain reaction (PCR)

RNA from tumor tissue samples was extracted using the mir-Vana™ Kit according to the instructions of the manufacturer. Alternatively, total RNA from primary cultures of GBMs and tumoral glial cell lines was prepared with Tri-Reagent as specified by the manufacturer. In all cases, RNA samples were treated with DNase, using Turbo DNA-free kit. RNA purity was assessed by spectrophotometric measurement of the $\text{OD}_{260}/\text{OD}_{280}$ and $\text{OD}_{260}/\text{OD}_{230}$ ratios with acceptable values falling between 1.8 and 2.1. RNA integrity was verified by Bioanalyzer (Agilent Technologies). RNA (1 μg) was incubated at 70°C for 10 min in the presence of 50 ng/mL of random hexamers. After cooling at 4°C , RNA was reverse transcribed into cDNA in a reaction buffer containing 10 mM DTT, 1 mM deoxynucleotide triphosphates, 1.2 U/ μL RNase inhibitor, and 10 U/ μL of MMLV-RT. The reaction proceeded at 37°C for 1.5 h, then at 70°C for 10 min. SYBR Green real-time reverse transcription (RT)-PCR for the genes of interest was done using specific primers. For normalization, RT-PCR was performed using eukaryotic 18s rRNA endogenous control primers. Oligonucleotide sequences used in this study are reported in Table 4.

Table 3. Clinical and histopathological informations in 34 gliomas analyzed

ID	Age	M/F	Pathology	Treatment
GBM.1	72	H	GBM	STUPP
GBM.2	77	H	GBM	BCNU+TMZ
GBM.3	81	H	GBM	STUPP
GBM.4	58	H	GBM	STUPP
GBM.5	59	H	GBM	STUPP
GBM.6	65	H	GBM	STUPP
GBM.7	58	H	GBM	STUPP
GBM.8	76	F	GBM	BCNU+TMZ
GBM.9	42	F	GBM	STUPP
GBM.10	69	H	GBM	STUPP
GBM.11	65	H	GBM	STUPP
GBM.12	56	H	GBM	STUPP
GBM.13	53	H	GBM	STUPP
GBM.14	69	H	GBM	STUPP
GBM.15	68	H	GBM	STUPP
GBM.16	50	F	GBM	STUPP
GBM.17	80	H	GBM	BCNU+TMZ
GBM.18	65	H	GBM	STUPP
GBM.19	58	H	GBM	STUPP
GBM.20	59	H	GBM	STUPP
GBM.21	58	H	GBM	STUPP
GBM.22	42	F	GBM	STUPP
GBM.23	65	H	GBM	STUPP
GBM.24	69	H	GBM	STUPP
ODG.25	27	H	oligodendroglioma	-
AO.26	53	F	anaplastic oligodendroglioma	TMZ
AA.27		F	anaplastic astrocytoma	-
OA.28	45	H	oligoastrocytoma	-
OA.29	49	F	oligoastrocytoma	TMZ
AA.30	43	H	anaplastic astrocytoma	RT
AA.31	64	H	anaplastic astrocytoma	VCR+RT
ODG.32	52	H	oligodendroglioma	RT
ODG.33	40	H	oligodendroglioma	-
ODG.34	41	H	oligodendroglioma	RT

RT, radiation therapy; TMZ, temozolomide; BCNU, bis-chloroethylnitrosourea; VCR, vincristine; STUPP, radiotherapy plus concomitant and adjuvant temozolomide.

Specific quantitative real-time PCR experiments for microRNAs were performed using TaqMan MicroRNA assay. A total of 40 ng of RNA was used to reverse transcribe specific miRNA of interest into cDNA using the TaqMan® miRNA reverse transcription kit. The reverse transcription primers for miR21 and

Table 4. Sequences of forward and reverse primers used to quantify human mRNAs

Gene	Primers sequences
YKL-40	Forward primer 5'-GGACCACAGG CCATCACAGT-3'
	Reverse primer 5'-CCAGCCTCAA CATGTACCCC-3'
Vimentin	Forward primer 5'-AAAGTGTGGC TGCCAAGAAC-3'
	Reverse primer 5'-AGCCTCAGAG AGGTCAGCAA-3'
MGMT	Forward primer 5'-GGGTCTGCAC GAAATAAAGC-3'
	Reverse primer 5'-CTCCGGACCT CCGAGAAC-3'
PDGFR α	Forward primer 5'-GGACTTACCC TGGAGAAGTG AA-3'
	Reverse primer 5'-CCAATTGAT GGATGGGACT-3'
TIMP1	Forward primer 5'-CTGTTGTTGC TGTGGCTGAT-3'
	Reverse primer 5'-AACTTGCCCC TGATGACG-3'
Bcl2 α	Forward primer 5'-GCACCTGCAC ACCTGGAT-3'
	Reverse primer 5'-AGGGCCAAAC TGAGCAGA-3'
Bax α	Forward primer 5'-ATGTTTTCTG ACGGCAACTT C-3'
	Reverse primer 5'-ATCAGTTCCG GCACCTTG-3'
18s	Forward primer 5'-CTACCACATC CAAGGAAGGC A-3'
	Reverse primer 5'-TTTTTCGTCA CTACCTCCCC G-3'
BCAN	Qantitect Primer Assay: Hs_BCAN_1_SG (Qiagen)

miR200a are hairpin primers specific for the mature miRNAs and do not bind to the miRNA precursor forms. Reverse transcription was followed by real-time PCR using miRNAs specific TaqMan® probe assays (miR21-5p, ID 000397; miR200a-3p, ID 002274; control snU6, ID 001973) on a Roche LightCycler 480. Expression data were normalized according to the expression of snU6. The relative mRNA or miRNA levels were calculated using the comparative Ct method ($\Delta\Delta C_t$).

MGMT promoter methylation analysis

Genomic DNA was prepared as described by the manufacturer then converted by using the EpiTect bisulfite kit. Methylation analysis was performed using MethyLight Technique as described previously.⁶⁷ Briefly, real-time, fluorescence-based polymerase chain reaction was performed using the LightCycler 480 (Roche Diagnostics). Bisulfite-converted genomic DNA was amplified using a set of primers and a fluorescent dye-labeled oligonucleotide probe, resulting in a semiquantitative methylation analysis. For the CpG islands of the 15 investigated gene regions (GENE), the primers and probes were designed specifically for methylated DNA. All primer and probe sequences have been published elsewhere, and the performance of the technique has been validated.^{67,68} The differences in amounts of input genomic DNA were normalized by the collagen Type II, $\alpha 1$ gene (Col2A1). SssI-treated genomic DNA was used as a standard.

Cell growth assay

Cells were plated in 24 well-plates (10000–20000 cells/well). Two days later, culture medium was renewed and cells received TMZ for 3 d, at the concentrations indicated. When O⁶-BG treatments were performed, cells were pre-treated with 50 μ M O⁶-BG for 4 h before the addition of TMZ. Control cells were treated in the presence of DMSO diluant. At the end of the treatment, cell proliferation was assessed with MTT. Briefly, 0.1 volume of MTT (5 mg/mL) was added in culture medium. After incubation at 37 °C for 1 h-2 h, culture medium was discarded and cells containing MTT formazan crystals were lysed in isopropanol. Optical density was measured at 570 nm, with background subtraction at 690 nm.

Cell transfection

The effect of miR21 and miR200a on chemosensitivity was evaluated by transfecting tumoral glial cells with miR200a precursor (pre-miR-200a) or antisense miR21 (anti-miR21) oligonucleotides (assay ID, PM10991 and AM10206, respectively), at 35 nM and 100 nM final concentrations, respectively. Cells were plated at 60000 cells per well in 12-well culture plates. Transfection was performed 24 h later in serum-free medium, using Oligofectamine transfection reagent and following the instructions of the manufacturers. In parallel, control cells were transfected with pre-miRNA or anti-miRNA negative controls. After 24 h, the medium was replaced with fresh medium containing 10% FBS. To evaluate the effects of the transfection on cell proliferation, cells were allowed to grow for additional 72 h in the absence or in the presence of TMZ as indicated. The expression of MGMT was evaluated 48 h post-transfection.

Statistical analysis

The Graph Prism version 6.0 software was used to generate statistical data. Statistical analysis of differences between miRNA levels in glioblastomas and lower grade brain tumors were evaluated using the non-parametric Mann–Whitney U-test between two groups. Correlations were calculated using Spearman correlation test. Significance was accepted at $P < 0.05$.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We are grateful to Dr Dominique Lombardo for helpful comments on the manuscript, and to Pr Dominique Figarella-Branger for histo-pathological examination and clinical informations regarding glioma samples. We thank Charles Prevot for the FACS analysis of cell cycle.

Supplemental Materials

Supplemental materials may be found here:
www.landesbioscience.com/journals/cbt/article/28920/

References

- Tate MC, Aghi MK. Biology of angiogenesis and invasion in glioma. *Neurotherapeutics* 2009; 6:447-57; PMID:19560735; <http://dx.doi.org/10.1016/j.neurt.2009.04.001>
- Bauchet L, Mathieu-Daudé H, Fabbro-Peray P, Rigau V, Fabbro M, Chinot O, Pallusseau L, Carmin C, Lainé K, Schlama A, et al.; Société Française de Neurochirurgie (SFNC); Club de Neuro-Oncologie of the Société Française de Neurochirurgie (CNO-SFNC); Société Française de Neuropathologie (SFNP); Association des Neuro-Oncologues d'Expression Française (ANOCEF). Oncological patterns of care and outcome for 952 patients with newly diagnosed glioblastoma in 2004. *Neuro Oncol* 2010; 12:725-35; PMID:20364023; <http://dx.doi.org/10.1093/neuonc/nuq030>
- Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJ, Janzer RC, Ludwin SK, Allgeier A, Fisher B, Belanger K, et al.; European Organisation for Research and Treatment of Cancer Brain Tumour and Radiation Oncology Groups; National Cancer Institute of Canada Clinical Trials Group. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol* 2009; 10:459-66; PMID:19269895; [http://dx.doi.org/10.1016/S1470-2045\(09\)70025-7](http://dx.doi.org/10.1016/S1470-2045(09)70025-7)
- Bocangel DB, Finkelstein S, Schold SC, Bhakat KK, Mitra S, Kokkinakis DM. Multifaceted resistance of gliomas to temozolomide. *Clin Cancer Res* 2002; 8:2725-34; PMID:12171906
- Oliva CR, Nozell SE, Diers A, McCluggage SG 3rd, Sarkaria JN, Markert JM, Darley-Usmar VM, Bailey SM, Gillespie GY, Landar A, et al. Acquisition of temozolomide chemoresistance in gliomas leads to remodeling of mitochondrial electron transport chain. *J Biol Chem* 2010; 285:39759-67; PMID:20870728; <http://dx.doi.org/10.1074/jbc.M110.147504>
- Le Calvé B, Rynkowski M, Le Mercier M, Bruyère C, Lonz C, Gras T, Haibe-Kains B, Bontempi G, Decaestecker C, Ruysschaert JM, et al. Long-term in vitro treatment of human glioblastoma cells with temozolomide increases resistance in vivo through up-regulation of GLUT transporter and aldo-keto reductase enzyme AKR1C expression. *Neoplasia* 2010; 12:727-39; PMID:20824049
- van Niftrik KA, van den Berg J, van der Meide WF, Ameziane N, Wedekind LE, Steenbergen RD, Leenstra S, Lafleur MV, Slotman BJ, Stalpers LJ, et al. Absence of the MGMT protein as well as methylation of the MGMT promoter predict the sensitivity for temozolomide. *Br J Cancer* 2010; 103:29-35; PMID:20517307; <http://dx.doi.org/10.1038/sj.bjc.6605712>
- Spiegel-Kreinecker S, Pirkner C, Filipits M, Lötsch D, Buchroithner J, Pichler J, Silye R, Weis S, Micksche M, Fischer J, et al. O6-Methylguanine DNA methyltransferase protein expression in tumor cells predicts outcome of temozolomide therapy in glioblastoma patients. *Neuro Oncol* 2010; 12:28-36; PMID:20150365; <http://dx.doi.org/10.1093/neuonc/nop003>
- Chinot OL, Barrié M, Fuentes S, Eudes N, Lancelot S, Metellus P, Muracciole X, Braguer D, Ouafik L, Martin PM, et al. Correlation between O6-methylguanine-DNA methyltransferase and survival in inoperable newly diagnosed glioblastoma patients treated with neoadjuvant temozolomide. *J Clin Oncol* 2007; 25:1470-5; PMID:17442989; <http://dx.doi.org/10.1200/JCO.2006.07.4807>
- Gaspar N, Marshall L, Perryman L, Bax DA, Little SE, Viana-Pereira M, Sharp SY, Vassal G, Pearson AD, Reis RM, et al. MGMT-independent temozolomide resistance in pediatric glioblastoma cells associated with a PI3-kinase-mediated HOX/stem cell gene signature. *Cancer Res* 2010; 70:9243-52; PMID:20935218; <http://dx.doi.org/10.1158/0008-5472.CAN-10-1250>
- Bobola MS, Kolstoe DD, Blank A, Silber JR. Minimally cytotoxic doses of temozolomide produce radiosensitization in human glioblastoma cells regardless of MGMT expression. *Mol Cancer Ther* 2010; 9:1208-18; PMID:20457618; <http://dx.doi.org/10.1158/1535-7163.MCT-10-0010>
- Zhang J, Stevens MF, Loughton CA, Madhusudan S, Bradshaw TD. Acquired resistance to temozolomide in glioma cell lines: molecular mechanisms and potential translational applications. *Oncology* 2010; 78:103-14; PMID:20357518; <http://dx.doi.org/10.1159/000306139>
- Jiang Z, Pore N, Cerniglia GJ, Mick R, Georgescu MM, Bernhard EJ, Hahn SM, Gupta AK, Maity A. Phosphatase and tensin homologue deficiency in glioblastoma confers resistance to radiation and temozolomide that is reversed by the protease inhibitor nelfinavir. *Cancer Res* 2007; 67:4467-73; PMID:17483362; <http://dx.doi.org/10.1158/0008-5472.CAN-06-3398>
- Dai C, Zhang B, Liu X, Ma S, Yang Y, Yao Y, Feng M, Bao X, Li G, Wang J, et al. Inhibition of PI3K/AKT/mTOR pathway enhances temozolomide-induced cytotoxicity in pituitary adenoma cell lines in vitro and xenografted pituitary adenoma in female nude mice. *Endocrinology* 2013; 154:1247-59; PMID:23384836; <http://dx.doi.org/10.1210/en.2012-1908>
- Cahill DP, Levine KK, Betensky RA, Codd PJ, Romany CA, Reavie LB, Barchelor TT, Futreal PA, Stratton MR, Curry WT, et al. Loss of the mismatch repair protein MSH6 in human glioblastomas is associated with tumor progression during temozolomide treatment. *Clin Cancer Res* 2007; 13:2038-45; PMID:17404084; <http://dx.doi.org/10.1158/1078-0432.CCR-06-2149>
- Happold C, Roth P, Wick W, Schmidt N, Florea AM, Silgner M, Reifenberger G, Weller M. Distinct molecular mechanisms of acquired resistance to temozolomide in glioblastoma cells. *J Neurochem* 2012; 122:444-55; PMID:22564186; <http://dx.doi.org/10.1111/j.1471-4159.2012.07781.x>
- Blough MD, Beauchamp DC, Westgate MR, Kelly JJ, Cairncross JG. Effect of aberrant p53 function on temozolomide sensitivity of glioma cell lines and brain tumor initiating cells from glioblastoma. *J Neurooncol* 2011; 102:1-7; PMID:20593219; <http://dx.doi.org/10.1007/s11060-010-0283-9>
- Chen CH, Chang YJ, Ku MSB, Chung KT, Yang JT. Enhancement of temozolomide-induced apoptosis by valproic acid in human glioma cell lines through redox regulation. *J Mol Med (Berl)* 2011; 89:303-15; PMID:21340685; <http://dx.doi.org/10.1007/s00109-010-0707-1>
- Momota H, Nerio E, Holland EC. Perifosine inhibits multiple signaling pathways in glial progenitors and cooperates with temozolomide to arrest cell proliferation in gliomas in vivo. *Cancer Res* 2005; 65:7429-35; PMID:16103096; <http://dx.doi.org/10.1158/0008-5472.CAN-05-1042>
- Zhang W, Zhang J, Yan W, You G, Bao Z, Li S, Kang C, Jiang C, You Y, Zhang Y, et al. Whole-genome microRNA expression profiling identifies a 5-microRNA signature as a prognostic biomarker in Chinese patients with primary glioblastoma multiforme. *Cancer* 2013; 119:814-24; PMID:22990979; <http://dx.doi.org/10.1002/cncr.27826>
- Ciafrè SA, Galardi S, Mangiola A, Ferracin M, Liu CG, Sabatino G, Negrini M, Maira G, Croce CM, Farace MG. Extensive modulation of a set of microRNAs in primary glioblastoma. *Biochem Biophys Res Commun* 2005; 334:1351-8; PMID:16039986; <http://dx.doi.org/10.1016/j.bbrc.2005.07.030>
- Niyazi M, Zehentmayr F, Niemöller OM, Eigenbrod S, Kretschmar H, Schulze-Osthoff K, Tonn JC, Atkinson M, Mörtl S, Belka C. MiRNA expression patterns predict survival in glioblastoma. *Radiat Oncol* 2011; 6:153-61; PMID:22074483; <http://dx.doi.org/10.1186/1748-717X-6-153>
- Wahid F, Shehzad A, Khan T, Kim YY. MicroRNAs: synthesis, mechanism, function, and recent clinical trials. *Biochim Biophys Acta* 2010; 1803:1231-43; PMID:20619301; <http://dx.doi.org/10.1016/j.bbamcr.2010.06.013>
- Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor suppressors. *Dev Biol* 2007; 302:1-12; PMID:16989803; <http://dx.doi.org/10.1016/j.jydbio.2006.08.028>
- DeSano JT, Xu L. MicroRNA regulation of cancer stem cells and therapeutic implications. *AAPS J* 2009; 11:682-92; PMID:19842044; <http://dx.doi.org/10.1208/s12248-009-9147-7>
- Hernando E. microRNAs and cancer: role in tumorigenesis, patient classification and therapy. *Clin Transl Oncol* 2007; 9:155-60; PMID:17403626; <http://dx.doi.org/10.1007/s12094-007-0029-0>
- Papagiannakopoulos T, Shapiro A, Kosik KS. MicroRNA-21 targets a network of key tumor-suppressive pathways in glioblastoma cells. *Cancer Res* 2008; 68:8164-72; PMID:18829576; <http://dx.doi.org/10.1158/0008-5472.CAN-08-1305>
- Conti A, Aguenouz M, La Torre D, Tomasello C, Cardali S, Angileri FF, Maio F, Cama A, Germanò A, Vita G, et al. miR-21 and 221 upregulation and miR-181b downregulation in human grade II-IV astrocytic tumors. *J Neurooncol* 2009; 93:325-32; PMID:19159078; <http://dx.doi.org/10.1007/s11060-009-9797-4>
- Hermansen SK, Dahlrot RH, Nielsen BS, Hansen S, Kristensen BW. MiR-21 expression in the tumor cell compartment holds unfavorable prognostic value in gliomas. *J Neurooncol* 2013; 111:71-81; PMID:23104517; <http://dx.doi.org/10.1007/s11060-012-0992-3>
- Lakomy R, Sana J, Hankeova S, Fadrus P, Kren L, Lzicarova E, Svoboda M, Dolezelova H, Smrcka M, Vyzula R, et al. MiR-195, miR-196b, miR-181c, miR-21 expression levels and O-6-methylguanine-DNA methyltransferase methylation status are associated with clinical outcome in glioblastoma patients. *Cancer Sci* 2011; 102:2186-90; PMID:21895872; <http://dx.doi.org/10.1111/j.1349-7006.2011.02092.x>
- Rossi S, Shimizu M, Barbarotto E, Nicoloso MS, Dimitri F, Sampath D, Fabbri M, Lerner S, Barron LL, Rassenti LZ, et al. microRNA fingerprinting of CLL patients with chromosome 17p deletion identify a miR-21 score that stratifies early survival. *Blood* 2010; 116:945-52; PMID:20393129; <http://dx.doi.org/10.1182/blood-2010-01-263889>
- Gao W, Yu Y, Cao H, Shen H, Li X, Pan S, Shu Y. Deregulated expression of miR-21, miR-143 and miR-181a in non small cell lung cancer is related to clinicopathologic characteristics or patient prognosis. *Biomed Pharmacother* 2010; 64:399-408; PMID:20363096; <http://dx.doi.org/10.1016/j.biopha.2010.01.018>
- Hwang JH, Voortman J, Giovannetti E, Steinberg SM, Leon LG, Kim YT, Funel N, Park JK, Kim MA, Kang GH, et al. Identification of microRNA-21 as a biomarker for chemoresistance and clinical outcome following adjuvant therapy in resectable pancreatic cancer. *PLoS One* 2010; 5:e10630; PMID:20498843; <http://dx.doi.org/10.1371/journal.pone.0010630>

34. Burk U, Schubert J, Wellner U, Schmalhofer O, Vincan E, Spaderna S, Brabletz T. A reciprocal repression between ZEB1 and members of the miR-200 family promotes EMT and invasion in cancer cells. *EMBO Rep* 2008; 9:582-9; PMID:18483486; <http://dx.doi.org/10.1038/embor.2008.74>
35. Gregory PA, Bert AG, Paterson EL, Barry SC, Tsykin A, Farshid G, Vadas MA, Khew-Goodall Y, Goodall GJ. The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat Cell Biol* 2008; 10:593-601; PMID:18376396; <http://dx.doi.org/10.1038/ncb1722>
36. Leskelä S, Leandro-García LJ, Mendiola M, Barriuso J, Inglada-Pérez L, Muñoz I, Martínez-Delgado B, Redondo A, de Santiago X, Robledo M, et al. The miR-200 family controls β -tubulin III expression and is associated with paclitaxel-based treatment response and progression-free survival in ovarian cancer patients. *Endocr Relat Cancer* 2011; 18:85-95; PMID:21051560; <http://dx.doi.org/10.1677/ERC-10-0148>
37. Shimono Y, Zabala M, Cho RW, Lobo N, Dalerba P, Qian D, Diehn M, Liu H, Panula SP, Chiao E, et al. Downregulation of miRNA-200c links breast cancer stem cells with normal stem cells. *Cell* 2009; 138:592-603; PMID:19665978; <http://dx.doi.org/10.1016/j.cell.2009.07.011>
38. Wiklund ED, Bramsen JB, Hulf T, Dyrskjot L, Ramanathan R, Hansen TB, Villadsen SB, Gao S, Ostenfeld MS, Borre M, et al. Coordinated epigenetic repression of the miR-200 family and miR-205 in invasive bladder cancer. *Int J Cancer* 2011; 128:1327-34; PMID:20473948; <http://dx.doi.org/10.1002/ijc.25461>
39. Li Y, VandenBoom TG, Kong D, Wang Z, Ali S, Philip PA, Sarkar FH. Up-regulation of miR-200 and let-7 by natural agents leads to the reversal of epithelial-to-mesenchymal transition in gemcitabine-resistant pancreatic cancer cells. *Cancer Res* 2009; 69:6704-12; PMID:19654291; <http://dx.doi.org/10.1158/0008-5472.CAN-09-1298>
40. Adam L, Zhong M, Choi W, Qi W, Nicoloso M, Arora A, Calin G, Wang H, Siefker-Radtke A, McConkey D, et al. miR-200 expression regulates epithelial-to-mesenchymal transition in bladder cancer cells and reverses resistance to epidermal growth factor receptor therapy. *Clin Cancer Res* 2009; 15:5060-72; PMID:19671845; <http://dx.doi.org/10.1158/1078-0432.CCR-08-2245>
41. Mueller DW, Rehli M, Bosserhoff AK. miRNA expression profiling in melanocytes and melanoma cell lines reveals miRNAs associated with formation and progression of malignant melanoma. *J Invest Dermatol* 2009; 129:1740-51; PMID:19212343; <http://dx.doi.org/10.1038/jid.2008.452>
42. Schultz J, Lorenz P, Gross G, Ibrahim S, Kunz M. MicroRNA let-7b targets important cell cycle molecules in malignant melanoma cells and interferes with anchorage-independent growth. *Cell Res* 2008; 18:549-57; PMID:18379589; <http://dx.doi.org/10.1038/cr.2008.45>
43. Iorio MV, Visone R, Di Leva G, Donati V, Petrocca F, Casalini P, Taccioli C, Volinia S, Liu CG, Alder H, et al. MicroRNA signatures in human ovarian cancer. *Cancer Res* 2007; 67:8699-707; PMID:17875710; <http://dx.doi.org/10.1158/0008-5472.CAN-07-1936>
44. Xi Y, Formentini A, Chien M, Weir DB, Russo JJ, Ju J, Kornmann M, Ju J. Prognostic values of microRNAs in colorectal cancer. *Biomark Insights* 2006; 2:113-21; PMID:18079988
45. Saydam O, Shen Y, Würdinger T, Senol O, Boke E, James MF, Tannous BA, Stemmer-Rachamimov AO, Yi M, Stephens RM, et al. Downregulated microRNA-200a in meningiomas promotes tumor growth by reducing E-cadherin and activating the Wnt/beta-catenin signaling pathway. *Mol Cell Biol* 2009; 29:5923-40; PMID:19703993; <http://dx.doi.org/10.1128/MCB.00332-09>
46. Hayashi T, Adachi K, Ohba S, Hirose Y. The Cdk inhibitor flavopiridol enhances temozolomide-induced cytotoxicity in human glioma cells. *J Neurooncol* 2013; 115:169-78; PMID:23943501; <http://dx.doi.org/10.1007/s11060-013-1220-5>
47. Shen W, Hu J-A, Zheng J-S. Mechanism of temozolomide-induced antitumor effects on glioma cells. *J Int Med Res* 2014; 42:164-72; PMID:24326954; <http://dx.doi.org/10.1177/0300060513501753>
48. Filippi-Chiela EC, Thomé MP, Bueno e Silva MM, Pelegrini AL, Ledur PF, Garicochea B, Zamin LL, Lenz G. Resveratrol abrogates the temozolomide-induced G2 arrest leading to mitotic catastrophe and reinforces the temozolomide-induced senescence in glioma cells. *BMC Cancer* 2013; 13:147-56; PMID:23522185; <http://dx.doi.org/10.1186/1471-2407-13-147>
49. Rickman DS, Bobek MP, Misk DE, Kuick R, Blaivas M, Kurnit DM, Taylor J, Hanash SM. Distinctive molecular profiles of high-grade and low-grade gliomas based on oligonucleotide microarray analysis. *Cancer Res* 2001; 61:6885-91; PMID:11559565
50. van den Boom J, Wolter M, Kuick R, Misk DE, Youkilis AS, Wechsler DS, Sommer C, Reifemberger G, Hanash SM. Characterization of gene expression profiles associated with glioma progression using oligonucleotide-based microarray analysis and real-time reverse transcription-polymerase chain reaction. *Am J Pathol* 2003; 163:1033-43; PMID:12937144; [http://dx.doi.org/10.1016/S0002-9440\(10\)63463-3](http://dx.doi.org/10.1016/S0002-9440(10)63463-3)
51. Healy NA, Heneghan HM, Miller N, Osborne CK, Schiff R, Kerin MJ. Systemic miRNAs as potential biomarkers for malignancy. *Int J Cancer* 2012; 131:2215-22; PMID:22618667; <http://dx.doi.org/10.1002/ijc.27642>
52. Nair VS, Maeda LS, Ioannidis JP. Clinical outcome prediction by microRNAs in human cancer: a systematic review. *J Natl Cancer Inst* 2012; 104:528-40; PMID:22395642; <http://dx.doi.org/10.1093/jnci/djs027>
53. Qiu S, Lin S, Hu D, Feng Y, Tan Y, Peng Y. Interactions of miR-323/miR-329 and miR-130a/miR155/miR-210 as prognostic indicators for clinical outcome of glioblastoma patients. *J Transl Med* 2013; 11:10; PMID:23281771; <http://dx.doi.org/10.1186/1479-5876-11-10>
54. Fu J, Rodova M, Nanta R, Meeker D, Van Veldhuizen PJ, Srivastava RK, Shankar S. NPV-LDE-225 (Erisomidegib) inhibits epithelial mesenchymal transition and self-renewal of glioblastoma initiating cells by regulating miR-21, miR-128, and miR-200. *Neuro Oncol* 2013; 15:691-706; PMID:23482671; <http://dx.doi.org/10.1093/neuonc/not011>
55. Metellus P, Coulilaly B, Nanni I, Fina F, Eudes N, Giorgi R, Barrie M, Chinot O, Fuentes S, Dufour H, et al. Prognostic impact of O6-methylguanine-DNA methyltransferase silencing in patients with recurrent glioblastoma multiforme who undergo surgery and carmustine wafer implantation. *Cancer* 2009; 115:4783-94; PMID:19637364; <http://dx.doi.org/10.1002/cncr.24546>
56. Haenisch S, Cascorbi I. miRNAs as mediators of drug resistance. *Epigenomics* 2012; 4:369-81; PMID:22920178; <http://dx.doi.org/10.2217/epi.12.39>
57. Ren Y, Zhou X, Mei M, Yuan XB, Han L, Wang GX, Jia ZF, Xu P, Pu PY, Kang CS. MicroRNA-21 inhibitor sensitizes human glioblastoma cells U251 (PTEN-mutant) and LN229 (PTEN-wild type) to taxol. *BMC Cancer* 2010; 10:27-34; PMID:20113523; <http://dx.doi.org/10.1186/1471-2407-10-27>
58. Corsten MF, Miranda R, Kasmieh R, Krichevsky AM, Weissleder R, Shah K. MicroRNA-21 knockdown disrupts glioma growth in vivo and displays synergistic cytotoxicity with neural precursor cell delivered S-TRAIL in human gliomas. *Cancer Res* 2007; 67:8994-9000; PMID:17908999; <http://dx.doi.org/10.1158/0008-5472.CAN-07-1045>
59. Shi L, Chen J, Yang J, Pan T, Zhang S, Wang Z. MiR-21 protected human glioblastoma U87MG cells from chemotherapeutic drug temozolomide induced apoptosis by decreasing Bax/Bcl-2 ratio and caspase-3 activity. *Brain Res* 2010; 1352:255-64; PMID:20633539; <http://dx.doi.org/10.1016/j.brainres.2010.07.009>
60. Li Y, Zhao S, Zhen Y, Li Q, Teng L, Asai A, Kawamoto K. A miR-21 inhibitor enhances apoptosis and reduces G(2)-M accumulation induced by ionizing radiation in human glioblastoma U251 cells. *Brain Tumor Pathol* 2011; 28:209-14; PMID:21618027; <http://dx.doi.org/10.1007/s10014-011-0037-1>
61. Wong ST, Zhang XQ, Zhuang JT, Chan HL, Li CH, Leung GK. MicroRNA-21 inhibition enhances in vitro chemosensitivity of temozolomide-resistant glioblastoma cells. *Anticancer Res* 2012; 32:2835-41; PMID:22753745
62. Ali S, Ahmad A, Banerjee S, Padhye S, Dominiak K, Schaffert JM, Wang Z, Philip PA, Sarkar FH. Gemcitabine sensitivity can be induced in pancreatic cancer cells through modulation of miR-200 and miR-21 expression by curcumin or its analogue CDF. *Cancer Res* 2010; 70:3606-17; PMID:20388782; <http://dx.doi.org/10.1158/0008-5472.CAN-09-4598>
63. Noonan EM, Shah D, Yaffe MB, Lauffenburger DA, Samson LD. O6-Methylguanine DNA lesions induce an intra-S-phase arrest from which cells exit into apoptosis governed by early and late multi-pathway signaling network activation. *Integr Biol (Camb)* 2012; 4:1237-55; PMID:22892544; <http://dx.doi.org/10.1039/c2ib20091k>
64. Roos WP, Kaina B. DNA damage-induced cell death: from specific DNA lesions to the DNA damage response and apoptosis. *Cancer Lett* 2013; 332:237-48; PMID:22261329; <http://dx.doi.org/10.1016/j.canlet.2012.01.007>
65. Kim T, Veronesi A, Pichiorri F, Lee TJ, Jeon YJ, Volinia S, Pineau P, Marchio A, Palatini J, Suh SS, et al. p53 regulates epithelial-mesenchymal transition through microRNAs targeting ZEB1 and ZEB2. *J Exp Med* 2011; 208:875-83; PMID:21518799; <http://dx.doi.org/10.1084/jem.20110235>
66. Ouafik L, Berenguer-Daize C, Berthois Y. Adrenomedullin promotes cell cycle transit and up-regulates cyclin D1 protein level in human glioblastoma cells through the activation of c-Jun/JNK/AP-1 signal transduction pathway. *Cell Signal* 2009; 21:597-608; PMID:19166930; <http://dx.doi.org/10.1016/j.cellsig.2009.01.001>
67. Ogino S, Kawasaki T, Brahmandam M, Cantor M, Kirkner GJ, Spiegelman D, Makrigiorgos GM, Weisenberger DJ, Laird PW, Loda M, et al. Precision and performance characteristics of bisulfite conversion and real-time PCR (MethylLight) for quantitative DNA methylation analysis. *J Mol Diagn* 2006; 8:209-17; PMID:16645207; <http://dx.doi.org/10.2353/jmoldx.2006.050135>
68. Widschwendter M, Siegmund KD, Müller HM, Fiegl H, Marth C, Müller-Holzner E, Jones PA, Laird PW. Association of breast cancer DNA methylation profiles with hormone receptor status and response to tamoxifen. *Cancer Res* 2004; 64:3807-13; PMID:15172987; <http://dx.doi.org/10.1158/0008-5472.CAN-03-3852>